arXiv:1606.07159v1 [math.SP] 23 Jun 2016

SPECTRAL INCLUSION FOR C_0-SEMIGROUPS DRAZIN INVERTIBLE AND QUASI-FREDHOLM OPERATORS

A. TAJMOUATI, M. AMOUCH, M. R. F.Z. ALHOMIDI

Abstract. Let $(T(t))_{t\geq 0}$ be a C_0 semigroups and A be its infinitesimal generator. In this work, we prove that the spectral inclusion for $(T(t))_{t\geq 0}$ remains true for the Drazin invertible and Quasi-Fredholm spectra. Also, we will give conditions under which facts A is quasi-Fredholm, A is Drazin invertible and A is B-Fredholm are equivalent.

1. Introduction and preliminaries.

Let X a Banach space and $\mathcal{B}(X)$ the Banach algebra of all bounded linear operators on X. for $T \in \mathcal{B}(X)$, by T^*, $N(T)$, $R(T)$, $R^\infty(T) = \bigcap_{n\geq 0} R(T^n)$, $N^\infty(T) = \bigcup_{n\geq 0} N(T^n)$, $\rho(T)$ and $\sigma(T)$, we denote, respectively the adjoint, the null space, the range, the hyper-range, the hyper-kernel, the resolvent set and the spectrum of T.

Let $T \in B(X)$, the ascent $a(T)$ and the descent $d(T)$ of T are defined in [15] by $a(T) = \inf\{n \in \mathbb{N} : N(T^n) = N(T^{n+1})\}$ and $d(T) = \inf\{n \in \mathbb{N} : R(T^n) = R(T^n + 1)\}$, respectively.

The operator T is said to be Drazin invertible if $d(T) < \infty$ and $a(T) < \infty$. It is well known that T is Drazin invertible if and only if $T = T_1 \oplus T_2$ where T_1 is invertible and T_2 is nilpotent, see [9, Corollary 2.2]. This is equivalent to the fact that, there exists an integer n such that the space $R(T^n)$ is closed and the restriction of T of $R(T^n)$ viewed as a map from $R(T^n)$ into $R(T^n)$ is invertible, see [2, Theorem 2.5]. The Drazin spectrum of T is defined by

$$\sigma_D(T) = \{\lambda \in \mathbb{C} : \lambda - T \text{ is not Drazin invertible}\}.$$

Similarly, from [7], T is left (respect right) generalized Drazin invertible if and only if $T = T_1 \oplus T_2$ such that T_1 is left (respect right) invertible and T_2 is quasi-nilpotent.

Recall that T is said to be semi-regular or Kato operator, if $R(T)$ is closed and $N(T) \subset R^\infty(T)$, see for example [1]. In addition, T is said to be pseudo-Fredholm operator if there exist two closed T-invariant subspaces M and N such that $X = M \oplus N$ and $T = T_{|M} \oplus T_{|N}$ with $T_{|M}$ is semi-regular and $T_{|N}$ is nilpotent. This is equivalent to the fact that there exists an integer n such that $R(T^n)$ is closed and the restriction of T of $R(T^n)$ viewed as a map from $R(T^n)$ into $R(T^n)$ is semi-regular. The quasi-Fredholm spectrum is defined by

$$\sigma_{QF}(T) = \{\lambda \in \mathbb{C} : \lambda - T \text{ is not quasi-Fredholm}\},$$

see [7, 10, 11] for more information.

2000 *Mathematics Subject Classification.* 47A16, 47D06, 47D03.

Key words and phrases. Banach space operators; C_0 semigroups; Spectral inclusion ; Drazin invertible operator; Quasi-Fredholm operator.

Similarly, from [3] T is said to be a B-Fredholm operator, if there exists an integer n such that the space $R(T^n)$ is closed and the restriction of T of $R(T^n)$ viewed as a map from $R(T^n)$ into $R(T^n)$ is Fredholm.

Let $T \in B(X)$ and $x \in X$, the local resolvent of T at x noted $\rho_T(x)$ is defined as the union of all open subset U of $\mathbb{C}$ for which there is an analytic function $f : U \to X$ such that the equation $(T-\mu I)f(\mu) = x$ holds for all $\mu \in U$. The local spectrum $\sigma_T(x)$ of T at x is defined by $\sigma_T(x) = \mathbb{C} \setminus \rho_T(x)$. Evidently $\rho_T(x)$ is an open subset of $\mathbb{C}$ and $\sigma_T(x)$ is closed. If $f(z) = \sum_{i=0}^{\infty} x_i(z-\mu)^i$ (in a neighborhood of μ), then $\mu \in \rho_T(x)$ if and only if there exists a sequence $(x_i)_{i\geq 0} \subseteq X$, $x_0 = x$, $(T-\mu)x_{i+1} = x_i$, and $\sup_i ||x_i||^{\frac{1}{i}} < \infty$, see [8].

Let $\mathcal{T} = (T(t))_{t\geq 0}$ be a strongly continuous semigroup (C_0 semigroup in short) with infinitesimal generator A on X. We will denote the type (growth bound) of $\mathcal{T}$ by ω_0:

$$\omega_0 := \lim_{t\to\infty} \frac{\ln \|T't)\|}{t}$$

$$= \inf\big\{\omega \in \mathbb{R} \ : \ \text{there existe M such that } \|T(t)\| \leq Me^{\omega t}, t \geq 0\big\},$$

see [4, 5, 13] for more information. Also, in [4, 5, 13] the authors showed that

$$e^{t\nu(A)} \subseteq \nu(T(t)) \subseteq e^{t\nu(A)} \bigcup \{0\}$$

where $\nu(.) \in \{\sigma_p(.), \sigma_{ap}(.), \sigma_r(.)\}$ is the point spectrum, approximative spectrum or residual spectrum.
The semigroup $T(t)$ is called differentiable for $t > t_0$ if for every $x \in X$, $t \to T(t)x$ is differentiable for $t > t_0$. $T(t)$ is called differentiable if it is differentiable for $t > 0$. If $B(\lambda, t)x = \int_0^t e^{\lambda(t-s)}T(s)xds$, then $B(\lambda, t)x$ is differentiable in t with $B^{'}(\lambda, t)x = T(t)x + \lambda B(\lambda, t)x$ and $B^{'}_{\lambda}(t)$ is a bounded linear operator in X, see [5] and [13].

Spectral inclusions for various reduced spectra of a C_0 semigroup was studied by authors in [5] and [4] for point spectrum, approximative spectrum and residual spectrum. Also, the spectral equality for a C_0 semigroup was studied by authors in [14] for semi-regular, essentially semi-regular and semi-Fredholm spectrum, respectively. In this work, we will continue in this direction, we will prove that the spectral inclusion for Drazin and quasi Fredholm spectra. Also, we will show that if $(T(t))_{t\geq 0}$ is a C_0 semigroup with infinitesimal generator A such that the equality $\lim_{t\to\infty} \frac{1}{t^n}\|T(t)\| = 0$ holds for some $n \in \mathbb{N}$, then the infinitesimal generator A is quasi-Fredholm if and only if it is Drazin invertible if and only if it is B-Fredholm.

2. Main results

We start by given the following two lemmas which are proved in [5]. They will be used to prove our main result.

Lemma 2.1. [5] *Let $(A, D(A))$ be the infinitesimal generator of a strongly continuous semigroup $(T(t))_{t\geq 0}$ and $B(\lambda, t) = \int_0^t e^{\lambda(t-s)}T(s)xds$ is a bounded operator from X to $D(A)$. Then, for every $\lambda \in \mathbb{C}$, $t > 0$ and $n \in \mathbb{N}$, the following statements hold:*

(1) $(a) : (e^{\lambda t} - T(t))^n(x) = (\lambda - A)^n B(\lambda, t)^n(x),\ \lambda \in \mathbb{C}, x \in X;$
$(b) : (e^{\lambda t} - T(t))^n(x) = B(\lambda, t)^n(\lambda - A)^n(x),\ \lambda \in \mathbb{C}, x \in D(A).$
(2) $R(e^{\lambda t} - T(t))^n \subseteq R(\lambda - A)^n.$
(3) $N(\lambda - A)^n \subseteq N(e^{\lambda t} - T(t))^n.$

Lemma 2.2. [5] *Let $(A, D(A))$ be the infinitesimal generator of a strongly continuous semigroup $(T(t))_{t\geq 0}$ on X and assume that the restricted semigroup $(T(t)_|)_{t\geq 0}$ is strongly continuous on some $(T(t))_{t\geq 0}$-invariant Banach space $Y \hookrightarrow X$. Then the infinitesimal generator of $(T(t)|)_{t\geq 0}$ is the part $(A_|, D(A_|))$ of A in Y.*

The following two lemmas which are proved in [14] will be used in the sequel.

Lemma 2.3. [14, Lemma 3.1] *Let A the infinitesimal generator of C_0-semigroup $T(t)_{t\geq 0}$ and $B(\lambda; t) = \int_0^t e^{\lambda(t-s)}T(s)ds$ is a linear bounded operator on X. Then there exist C and D tow operator such that $(\lambda - A), B(\lambda, t), C, D$ are mutually commuting operators, for all $x \in D(A)$ and $C(\lambda - A) + DB(\lambda, t) = I$, for $t > 0$.*

Lemma 2.4. [14, Lemma 3.2] *Let $(\lambda - A), B(\lambda, t),\ C,\ D$ be mutually commuting operators in $D(A)$ such that $C(\lambda - A) + DB(\lambda, t) = I$, $t > 0$. Then we have:*

(1) *For every positive integer n there are $C_n, D_n \in D(A)$ such that $(\lambda - A)^n, B^n(\lambda, t), C_n, D_n$ are mutually commuting and*

$$(\lambda - A)^n C_n + B^n(\lambda, t)D_n = I.$$

(2) *For every positive integer n, $R(e^{\lambda t} - T(t))^n = R(\lambda - A)^n \bigcap R(B^n(\lambda, t))$ and $N((e^{\lambda t} - T(t))^n) = N((\lambda - A)^n) + N(B^n(\lambda, t))$. Further $R^\infty(e^{\lambda t} - T(t)) = R^\infty(\lambda - A) \bigcap R^\infty(B(\lambda, t))$ and $N^\infty(e^{\lambda t} - T(t)) = N^\infty(\lambda - A) + N^\infty(B(\lambda, t))$.*
(3) *$N^\infty(\lambda - A) \subset R^\infty(B(\lambda, t))$ and $N^\infty(B(\lambda, t)) \subset R^\infty(\lambda - A)$.*

Now, we give some spectral results for differentiable C_0 semigroup.

Lemma 2.5. *Let $(A, D(A))$ be the infinitesimal generator of a strongly continuous semigroup $(T(t))_{t\geq 0}$, $B(\lambda, t) = \int_0^t e^{\lambda(t-s)}T(s)xds$ is a bounded operator from X to $D(A)$. If $T(t)$ is differentiable for $t > t_0$, then for every $\lambda \in \mathbb{C}$, $t > t_0$ and $n \in \mathbb{N}$, the following statements hold:*

(1) $(a) : (\lambda e^{\lambda t} - AT(t))^n(x) = (\lambda - A)^n B(\lambda, t)^{'n}(x),\ \lambda \in \mathbb{C}, x \in X;$
$(b) : (\lambda e^{\lambda t} - T(t))^n(x) = B(\lambda, t)^{'n}(\lambda - A)^n(x),\ \lambda \in \mathbb{C}, x \in D(A).$
(2) $R(\lambda e^{\lambda t} - AT(t))^n \subseteq R(\lambda - A)^n.$
(3) $N(\lambda - A)^n \subseteq N(\lambda e^{\lambda t} - AT(t))^n.$

Proof. Assuming now that $t > t_0$ and differentiating (a) and (b) in (1) of Lemma 2.1 with respect to t and $n = 1$, we obtain

$$\begin{aligned} \lambda e^{\lambda t}x - AT(t)x &= (\lambda I - A)B^{'}(\lambda, t)x \text{ for every } x \in X; \\ \lambda e^{\lambda t}x - AT(t)x &= B^{'}(\lambda, t)x(\lambda I - A) \text{ for every } x \in D(A). \end{aligned}$$

This gives (a) and (b) for $n = 1$. By induction, we obtain (a) and (b) for all $n \in \mathbb{N}$. The rest of Lemma follows from (1). □

Proposition 2.1. *Let $T(t)_{t>0}$ be a C_0 semigroup and let A be its infinitesimal generator. If $T(t)$ is differentiable for $t > t_0$ and $\lambda \in \sigma_A(x)$ for $x \in X$, then*

$$\lambda e^{\lambda t} \in \sigma_{AT(t)}(x).$$

Proof. Let $t > t_0$ be fixed and suppose that $\lambda e^{\lambda t} \notin \sigma_{AT(t_0)}(x)$, then there exist a sequence $(x_i)_{i\in\mathbb{N}}$ of X such that $x_0 = x$, $(\lambda e^{\lambda t} - AT(t))x_i = x_{i-1}$ and $\sup_i \|x_i\|^{\frac{1}{i}} < \infty$. We put $y_i = B^{'i}(\lambda, t)x_i$, as $B^{'}(\lambda, t)$ is a bounded linear operator in X, we have $y_0 = x_0 = x,\ y_0 \in D(A)$.

$$\begin{aligned}(\lambda - A)y_i &= (\lambda - A)B^{'}(\lambda, t)x_i B^{'(i-1)}(\lambda, t)x_i \\ &= (\lambda e^{\lambda t} - AT(t))B^{'(i-1)}(\lambda, t)x_i \\ &= B^{'(i-1)}(\lambda, t)(\lambda e^{\lambda t} - AT(t))x_i \\ &= B^{'(i-1)}(\lambda, t)x_{i-1} \\ &= y_{i-1}\end{aligned}$$

Therefor $(\lambda - A)y_i = y_{i-1}$. On the other hand

$$\|y_i\| = \|B^{'i}(\lambda, t)x_i\| < \|B^{'i}(\lambda, t)\|\|x_i\| < M^i\|x_i\|,$$

then

$$\sup_i \|y_i\|^{\frac{1}{i}} < \sup_i M\|x_i\|^{\frac{1}{i}} < \infty.$$

So that $\lambda \notin \sigma_A(x)$ □

Denote by $\sigma_{su}(A)$ the subjectivity spectrum of A. It is known that $\bigcup_{x\in X} \sigma_A(x) = \sigma_{su}(A)$ for a closed operator A. Hence the following corollary holds.

Corollairy 2.1. *Let $T(t)_{t>0}$ be a C_0 semigroup and let A be its infinitesimal generator. If $T(t)$ is differentiable for $t > t_0$ and $\lambda \in \sigma_{su}(A)$, then $\lambda e^{\lambda t} \in \sigma_{su}(AT(t))$.*

Proposition 2.2. *Let $T(t)_{t>0}$ be a C_0 semigroup and let A be its infinitesimal generator. If $T(t)$ is differentiable for $t > t_0$ and $\lambda \in \sigma_{ap}(A)$, then $\lambda e^{\lambda t} \in \sigma_{ap}(AT(t))$.*

Proof. For $t > t_0$, since $\lambda e^{\lambda t} - AT(t) = (\lambda - A)B^{'}(\lambda, t)$ and $B^{'}(\lambda, t)$ is bounded linear operator, then $\lambda e^{\lambda t} - AT(t)$ is a bounded linear operator. It is easy to check that for $x \in D(A)$, $B^{'}(\lambda, t)Ax = AB^{'}(\lambda, t)x$. If $\lambda \in \sigma_{ap}(A)$, then there exists sequence $(x_n)_{n\in\mathbb{N}} \in D(A)$ satisfying $||x_n|| = 1$ and $||(\lambda - A)x_n|| \to 0$. From (1) of lemma 2.5, we obtain the result. □

By an outline of the proof of [6, Theorem 2.1], we obtain the following result.

Proposition 2.3. *Let $T(t)_{t>0}$ be a C_0 semigroup and let A be its infinitesimal generator. If $T(t)$ is differentiable for $t > t_0$ and $\lambda \in \nu(A)$, then $\lambda e^{\lambda t} \in \nu(AT(t))$. where $\nu(.) \in \{\sigma_\gamma(.), \sigma_\pi(.), \sigma_{\gamma e}(.)\}$ and $\sigma_\gamma(.), \sigma_\pi(.), \sigma_{\gamma e}(.)$ denote the regular spectrum, essential regular spectrum and left essential spectrum.*

In the next theorem we will prove that the spectral inclusion of C_0semi-groups remains true for the Drazin invertible and quasi-Fredholm spectra.

Theorem 2.1. *Let $(T(t))_{t\geq 0}$ a C_0-semigroup, with infinitesimal generator A. Then*

$$e^{t\sigma_D(A)} \subseteq \sigma_D(T(t)).$$

Proof. Let $t_0 > 0$ be fixed and suppose that $(e^{\lambda t_0} - T(t_0))$ is Drazin invertible for some $\lambda \in \mathbb{C} \setminus \{0\}$. Then $M := R(e^{\lambda t_0} - T(t_0))^n$ is closed and the restricted semigroup $(e^{\lambda t_0} - T(t_0)_{|M})$ is invertible. We show that $(\lambda - A)$ is Drazin invertible. To this end, in the first we show that $R(\lambda - A)^n$ is closed. Let $x \in \overline{R((\lambda - A)^n))}$, that is, there exist $u_k \in D(A^n)$, $k = 1, 2, ...$, such that $(\lambda - A)^n u_k \to x$, hence

$$(e^{\lambda t_0} - T(t_0))^n u_k := B(\lambda, t_0)^n (\lambda - A)^n u_k$$

by Lemma 2.1. Also,

$$B(\lambda, t_0)^n (\lambda - A)^n u_k \ \to B(\lambda, t_0)^n x.$$

Hence

$$(e^{\lambda t_0} - T(t_0))^n u_k \ \to B(\lambda, t_0)^n x.$$

Since $M := R(e^{\lambda t_0} - T(t_0))^n$ is closed, then

$$B(\lambda, t_0)^n x \in R(e^{\lambda t_0} - T(t_0))^n.$$

Hence there exists $u \in D(A^n)$ such that

$$B(\lambda, t_0)^n x = (e^{\lambda t_0} - T(t_0))^n u.$$

In the other hand, From Lemma 2.1, we have that

$$(e^{\lambda t_0} - T(t_0))^n u = B(\lambda, t_0)^n (\lambda - A) u,$$

hence

$$B(\lambda, t_0)^n x = B(\lambda, t_0)^n (\lambda - A) u.$$

This implies that,

$$x - (\lambda - A)^n u \in N(B(\lambda, t_0)) \subseteq R(\lambda - A)^n.$$

So $x \in R(\lambda - A)^n$, and hence $R(\lambda - A)^n$ is closed.
Now, let us to show that $(\lambda - A_{|R(\lambda - A)^n})$ is invertible. For this, as $(e^{\lambda t_0} - T(t_0)_{|M})$ is invertible, then $(e^{\lambda t_0} - T(t_0)_{|M})$ is bounded below and $R(e^{\lambda t_0} - T(t_0)_{|M}) = R(e^{\lambda t_0} - T(t_0))^{n+1}$ is onto. We show that $(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is bounded below. Since $(e^{\lambda t_0} - T(t_0)_{|M})$ is bounded below, then $(e^{\lambda t_0} - T(t_0)_{|M})$ is injective and $R(e^{\lambda t_0} - T(t_0)_{|M})$ is closed. We show that $(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is injective and $(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is closed. For all $x \in D(A)$ we have

$$\{0\} = N(e^{\lambda t_0} - T(t_0)_{|M \bigcap D(A)}) = N(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)}) + N(B(\lambda, t_0)) \bigcap R(B^n(\lambda, t_0)),$$

then $N(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)}) = \{0\}$, therefore $(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is injective. As $R(e^{\lambda t_0} - T(t_0))^{n+1}$ is closed, then $(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)}) = R(\lambda - A)^{n+1}$ is also closed. On the other hand, as $R(e^{\lambda t_0} - T(t_0)_{|M})$ is onto, we can easily verify that $R(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is onto. In fact one can verify that

$$R(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)}) = R(\lambda - A)^{n+1} = R(\lambda - A)^n.$$

We have $R(\lambda - A)^{n+1} \subset R(\lambda - A)^n$ and if $y \in R(\lambda - A)^n$, then there exist $x \in D(A^n)$ such that $y = R(\lambda - A)^n x$ and by (1) of Lemma 2.3, we have

$$\begin{aligned}(\lambda - A)^n x &= (\lambda - A)^n C_n (\lambda - A)^n x + D_n B^n(\lambda, t)(\lambda - A)^n x \\ &= (\lambda - A)^{n+1} C_n (\lambda - A)^{n-1} x + D^n (e^{\lambda t_0} - T(t_0))^n x\end{aligned}$$

and as $R(e^{\lambda t_0} - T(t_0) \mid_{(e^{\lambda t_0} - T(t_0))^n}) = R(e^{\lambda t_0} - T(t_0))^{n+1}$ is onto, then there exist $x' \in X$ such that $(e^{\lambda t_0} - T(t_0))^n x = (e^{\lambda t_0} - T(t_0))^{n+1} x' = (\lambda - A)^{n+1} B^{n+1}(\lambda, t) x'$, therefore $y \in R(\lambda - A)^n$ and then $R(\lambda - A_{|R(\lambda - A)^n \bigcap D(A)})$ is onto.

Finally, $(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)})$ is Drazin invertible.

□

Corollairy 2.2. *For the infinitesimal generator A of a strongly continuous semigroup $(T(t))_{t\geq 0}$ one has the spectral inclusion*

$$e^{t\sigma_\nu(A)} \subseteq \sigma_\nu(T(t))$$

where $\sigma_\nu(.)$ is the left Drazin and right Drazin spectra.

The following example shows that the inclusion in Theorem 2.1 is strict.

Example 1. *Let X be the Banach space of continuous functions on $[0,1]$ which are equal to zero at $x = 1$ with the supremum norm. Define*

$$(T(t)f)(x) := \begin{cases} f(x+t) & if x+t \leq 1; \\ 0 & if x+t > 1. \end{cases}$$

$T(t)$ is obviously a C_0 semigroup on X.. Its infinitesimal generator A is given on

$$D(A) = \{f : f \in C^1([0,1]) \cup X, f^{'} \in X\}$$

by

$$Af = f^{'} \text{ for } f \in D(A).$$

One checks easily that for every $\lambda \in \mathbb{C}$ and $g \in X$ the equation $\lambda f - f^{'} = g$ has a unique solution $f \in X$ given by

$$f(t) = \int_t^1 e^{\lambda(t-S)} g(s) ds.$$

Therefore $\sigma(A) = \emptyset$ see [13]*, hence $\sigma_D(A) = \emptyset$.*
On the other hand, $T(t)$ is a bounded linear operator for every $t \geq 0$. Suppose that $f \in X$ is not periodic, then we have $0 \in \sigma_D(T(t))$. Indeed, if $0 \notin \sigma_D(T(t))$, then there exist a finite integer $n \in \mathbb{N}$ such that $(T^n(t)f)(x) = (T^{n+1}(t)f)(x)$, this implies that $f(x+nt) = f(x+nt+t)$, therefore f is periodic, this is a contradiction. Hence the inclusion

$$e^{t\sigma_D(A)} \subseteq \sigma_D(T(t)).$$

is strict.

Theorem 2.2. *For the infinitesimal generator A of a strongly continuous semigroup $(T(t))_{t\geq 0}$, we have the following inclusion:*

$$e^{t\sigma_{QF}(A)} \subseteq \sigma_{QF}(T(t)).$$

Proof. Let $t_0 > 0$ be fixed and suppose that $(e^{\lambda t_0} - T(t_0))$ is quasi-Fredholm, for some $\lambda \in \mathbb{C}\backslash\{0\}$. Then $M := R(e^{\lambda t_0} - T(t_0))^n$ is closed and the restricted semigroup $(e^{\lambda t_0} - T(t_0)_{|M})$ is semi-regular.
We show that $(\lambda - A)$ is quasi-Fredholm, to this end we show that $R(\lambda - A)^n$ is closed and $(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)})$ is semi-regular. That is, $(R(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)})$ is closed and $N(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)}) \subseteq R^\infty(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)}))$.
As $M := R(e^{\lambda t_0} - T(t_0))^n$ is closed, then by the same argument as in the proof of Theorem 2.1, we conclude that $R(\lambda - A)^n$ is closed. Since $(e^{\lambda t_0} - T(t_0)_{|M})$ is semi-regular, then $R(e^{\lambda t_0} - T(t_0)_{|M}) = R(e^{\lambda t_0} - T(t_0))^{n+1}$ is closed, this implies

that $R(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)}) = R(\lambda - A)^{n+1}$ is closed. On the other hand, by Lemma 2.1 we have that

$$N(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)}) \subseteq N(e^{\lambda t_0} - T(t_0)_{|M \bigcap D(A)} \subseteq N(e^{\lambda t_0} - T(t_0)_{|M}).$$

Since $(e^{\lambda t_0} - T(t_0)_{|M})$ is semi-regular, then $N(e^{\lambda t_0} - T(t_0)_{|M}) \subseteq R^\infty(e^{\lambda t} - T(t_0)_{|M}) = R^\infty(\lambda - A_{|R(\lambda-A) \bigcap D(A)}) \bigcap R^\infty B(\lambda, t_0) \subset R^\infty(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)})$, hence $(\lambda - A_{|R(\lambda-A)^n \bigcap D(A)})$ is semi-regular. Finally, we conclude that $(\lambda - A)$ is quasi-Fredholm. □

Let A be the infinitesimal generator of a C_0 semigroup $(T(t))_{t\geq 0}$. In the following, we will give condition on $(T(t))_{t\geq 0}$ under which facts A is Drazin invertible, A is $B-$Fredholm and A is $Q-$Fredholm are equivalent.

Theorem 2.3. *Let A be the infinitesimal generator of a C_0 semigroup $(T(t))_{t\geq 0}$. If $\lim_{t\to\infty} \frac{1}{t^n}\|T(t)\| = 0$, for some $n \in \mathbb{N}$, the following assertions are equivalent:*

1. *A is quasi-Fredholm;*
2. *A is Drazin invertible;*
3. *A is B-Fredholm.*

Proof. (1) $\Rightarrow$ (2) :

Since A is quasi-Fredholm, then $R(A^n)$ closed and $A_{|(R(A^n) \bigcap D(A))}$ is semi-regular. Let $y \in N(A_{|(R(A^n) \bigcap D(A))})$, then there exists $x \in (R(A^n) \bigcap D(A^n))$ such that $y = A^n x$. We integer by parts in the following formula:

$$T(t)x - x = \int_0^t T(s)Axds,$$

we obtain that

$$T(t)x = x + tA + \frac{t^2}{2!}A^2 + \int_0^t \frac{(t-s)^2}{2!}T(s)A^3xds.$$

We repeat this operation for n times, we obtain that

$$T(t)x = \sum_{k=0}^{n-1} \frac{t^k}{k!}A^k x + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!}T(s)A^n xds.$$

Hence,

$$\begin{aligned} T(t)x &= \sum_{k=0}^{n-1} \frac{t^k}{k!}A^k x + y\int_0^t \frac{(t-s)^{n-1}}{(n-1)!}ds \\ &= \sum_{k=0}^{n-1} \frac{t^k}{k!}A^k x + \frac{t^n}{n!}y. \end{aligned}$$

As $\lim\limits_{t\to\infty} \dfrac{1}{t^n}\|T(t)\| = 0$, then $y = 0$, this implies that

$$N(A_{|(R(A^n) \bigcap D(A))}) = \{0\}.$$

On the other hand, let $(T(t)^{'})_{t\geq 0}$ with infinitesimal generator $A^{'}$ the adjoint semigroup of $(T(t))_{t\geq 0}$. Since $A_{|(R(A^n) \bigcap D(A))}$ is semi-regular, then $A^{'}_{|(R(A'^n) \bigcap D(A'))}$ is also semi-regular, see [12, Proposition 1.6]. Using the following formula

$$T(t)^{'}x^{'} - x^{'} = weak^* \int_0^t T(s)^{'}A^{'}x^{'}ds,\ \forall x^{'} \in (R(A^{'n}) \bigcap D(A^{'})),\ \forall t \geq 0$$

which is proved in [16, Proposition 1.2.2] and by the same argument as above, we get that $N(A'_{|(R(A'^n)\bigcap D(A'))}) = \{0\}$. This is equivalent to the fact that

$$\overline{R(A_{|(R(A^n)\bigcap D(A))})} = (R(A^n)\bigcap D(A)).$$

Hence $R(A_{|(R(A^n)\bigcap D(A))}) = (R(A^n)\bigcap D(A))$, since $(R(A^n)\bigcap D(A))$ is closed. From this it follows that $A_{|(R(A^n)\bigcap D(A))}$ is surjective and hence it is invertible. Finally, A is Drazin invertible.
(2) $\Rightarrow$ (1): is clear.
(1), (2) and (3) are equivalent, since the class of Drazin invertible operator is a subclass of B-Fredholm operator and the class of B-Fredholm operator is a subclass of Quasi-Fredholm operator.

□

References

[1] P. Aiena, *Fredholm and local spectral theory with applications to multipliers,* Kluwer Academic Publishers.
[2] P.Aiena, Maria T. Biondi and C. Carbintero, *On Drazin Invertibility* , Amercivan. Mathemathecal. Society, (2008) 2839-2848.
[3] M. Berkani, *On a class of quasi-Fredholm operators*, Integral Equations Operator theory, **34** (1999), 244-603.
[4] Ph. Clment, H. J. A. M. Heijmans, S. Angenent, C. J. van Duijn, and B. de Pagter, *One-Parameter Semigroups*, Centre for Mathematics and Computer Science, (1987).
[5] K-J. Engel and R .Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag New York, (2000).
[6] A. Elkoutri and M. A. Taoudi, *Spectral Inclusions And Stability Results For Strongly Continuous Semigroups*, IJMMS **37**(2003) 2379-2387.
[7] K. M. Hocinea, M. Benharratb. and B. Messirdia, *Left and right generalized Drazin invertible operators*, Linear and Multilinear Algebra,(2014).
[8] K.B.Laursen, M.M.Neumann, *An introduction to Local Spectral Theory,* London Mathematical Society Monograph, New series, Vol. 20, Clarendon Press, Oxford, 2000
[9] D. C. Lay, *Spectral analysis using ascent, descent, nullity and defect*, Math. Ann. **184** (1970)197-214.
[10] M. Mbekhta, *Decomposition de Kato généralisée* , C. R. Acad. Sci. Paris 303 série I, No 20 (1990) 255-276.
[11] M. Mbekhta, *Opérateur pseudo -Fredholm I: Rsolvant généralisé,* J. Operator Theory, **24** (1990) 255-276.
[12] M. Mbekhta, *On the generalzed resolvent in Banach spaces*, J. Math. Anal. Appl. **189** (1995), 362-377.
[13] A.Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag New York, (1983).
[14] A. Tajmouati, M. Amouch, Z. Alhomidi, *Spectral Equality for C_0-Semigroups and Spectral inclusion of B-Fredholm*, arXiv:1601.04173.
[15] A.E. Taylor, D.C. Lay, *Introduction to Functional Analysis*, Wiley, New York-Chichester-Brisbane (1980).
[16] J.M.A.M.Van Neerven, *The Adjoint of a Semigroup of Linear Operator*, Lecture Notes in Mathe, **1529** (1992) Springer-Verlag.

A. TAJMOUATI and M. R. F. ALHOMIDI ZAKARIYA
Sidi Mohamed Ben Abdellah Univeristy Faculty of Sciences Dhar Al Mahraz Fez, Morocco.
E-mail address: abdelaziztajmouati@yahoo.fr
E-mail address: zakariya1978@yahoo.com

M. AMOUCH
Department of Mathematics University Chouaib Doukkali, Faculty of Sciences, Eljadida. 24000, Eljadida, Morocco.
E-mail address: mohamed.amouch@gmail.com